

Influence of Phytochemical Constituents of Garlic Extract (*Allium Sativum*) on the Treatment of Bacterial and Fungal Infections in *Clarias Gariepinus*

¹Sabo, Y. H, ²Sanchi, I.D and ³Alhassan, Y. J

¹Department of Water Resources, Aquaculture and Fisheries Technology,
Federal University of Technology, Minna Nigeria

²Department of Agricultural Economics and Extension, Federal University of Agriculture Zuru, Kebbi State,
Nigeria

³Department of Agricultural Economics and Extension, Federal University Wukari, Taraba State, Nigeria
yohannaj@fuwukari.edu.ng

Received: 7/3/2024 | Accepted: 23/3/2024 |

Abstract: Bacteria and fungi are among the most common pathogens that infect fish, leading to a significant economic lost in aquaculture. These microbial pathogens have also been reported to be of serious public health concern to humans. This study determined the influence of phytochemical constituent of garlic extract (*Allium sativum*) on the treatment of bacterial and fungal infection in *Clarias gariepinus*. Garlic bulbs were purchased from Kure market in Minna Niger State. The bulbs were peeled, washed under running tap water and pulverized using cold maceration method. Quantitative analysis was carried out to determine phytochemical compound (flavonoids, phenol, saponins, alkaloid and tannins) present in the extract. A total of thirty infected *Clarias gariepinus* were sampled from randomly selected fish farms. The total Bacteria, Coliform and Fungal count were determined and characteristically distinct colonies obtained were sub-cultured onto fresh agar plates repeatedly to obtain pure cultures which were then stored on appropriate agar slants for identification and further analysis. Bacterial species isolated were identified based on colony morphology and biochemical characteristics. The antimicrobial activity of *Allium sativum* was carried out using agar well diffusion method. Aqueous extraction of *Allium sativum* bulbs yielded 37.69g equivalent to 9.4% percentage yield. Quantitative phytochemical analysis revealed the presence of Phenols (205.96 mg/100g), Flavonoids (62.85 mg/100g), Saponin (55.22 mg/100g), Tannins (9.39 mg/100g) and Alkaloids (7.72 mg/100g). The total bacterial count of infected *Clarias gariepinus* samples range from 6×10^4 – 37×10^4 CFU/g with average count of 18.73×10^4 CFU/g while the coliform count ranged from 1×10^3 – 23×10^3 CFU/g and had an average count of 9.00×10^3 CFU/g. No fungal contamination was observed in this study. Bacteria isolated from infected *Clarias gariepinus* include members of the genus *Lactobacillus*, *Staphylococcus*, *Bacillus*, *Micrococcus*, *Streptococcus*, *Enterobacter* and *Escherichia*. Data indicated that the aqueous extracts of the *Allium sativum* bulbs had antimicrobial activity against suspected *Clarias gariepinus* pathogens. The antimicrobial activity was concentration dependent as inhibition zones increases with increasing concentration of the extract. With significantly highest antibacterial activity observed at 1000mg/ml. The results of this study indicate that garlic extract (*Allium sativum*) shows antibacterial potential in vitro. These shows that *Allium sativum* bulbs extracts are potentially effective as natural alternatives for the treatment of infections in *Clarias gariepinus*

Keywords: Influence, Phytochemical Constituents, Garlic Extract, Bacterial and Fungal Infection, *Clarias Gariepinus*

Published 30/3/2024

INTRODUCTION

Clarias gariepinus is a dominant freshwater fish and popular in commercial aquaculture due to its ability to grow rapidly and high tolerance to environmental conditions (Ayele, 2015). However, *C. gariepinus* has

been found to be susceptible to both microbial and parasitic infections particularly in intensive culture systems (Sudheesh et al., 2012; Opiyo et al., 2020). For decades' antibiotics have been used for the treatment of

bacterial infection in fish. The adverse effects associated with the use of antibiotics include drug residue, bioaccumulation and resistance of pathogens, which threaten human consumers (Ben et al., 2019). Hence, herbs are now being used as probiotics in preventing bacterial infections and are gaining success because they are cost effective, eco-friendly and have minimal side effects (Abd El-Hacket al., 2018). Herbs exhibit anti-microbial, anti-stress, appetite stimulation, immune stimulation, and aphrodisiac and antipathogenic effects which facilitate growth and maturation of cultured species (Nwabueze et al., 2020). *Allium sativum* is a pungent herb and has been reported to inhibit bacterial growth, promote fish growth and enhancement of blood parameters (Metwally, 2009; Alam et al., 2016). In aquaculture, *A. sativum* has been observed to promote growth, enhance immunity, stimulate appetite and strengthens the control of bacteria and fungi pathogens (Harris et al., 2001; Lee and Gao, 2012). *S. aureus* is one of the major bacterial agents causing food-borne diseases in humans worldwide and has been found on skin of healthy people and animals including fish (Fetsch and Jöhler, 2018). Economic Important of *Clarias gariepinus*

Fishes especially, catfish (*C. gariepinus*) are good sources of food for human beings. It is very rich in proteins and vitamins, especially, vitamin A (Retinol). They are source of animal protein. Fishes such as those in the class *Clariidae* are highly commercialized. Fishes have been known to feed on wide variety of things ranging from Sandy particles, phytoplankton, zooplanktons, leaves, roots, insects, insect larvae, worms, fishes etc. *C. gariepinus* is a benthic-pelagic fish which is known to have wide range of diet. Catfishes (*Clarias species*) are some of the most important fish species for aquaculture due to its high growth rate, significant tolerance to environment stress, reproduction in captivity, hardy to high density culture and its market demand. *C. gariepinus* especially is widely accepted by Nigerian consumers and was acknowledged that these bigger fish are sold for about twice the price of 30 days old fish. *C. gariepinus* has an average adult length of more than 1-meter long. These fish have slender bodies, a flat bony head, and a broad, terminal mouth with 4 pairs of barbells. They also have a large accessory breathing organ. They can weigh up to 29 kg or more.

Statement of the Research Problem

Fish disease is a substantial source of monetary loss to aqua culturists. Production costs are increased by fish disease outbreaks because it leads to investment lost in fish mortality, disease treatment, and decreased growth during convalescence. Fish are much less crowded in natural systems than in captivity. Parasites and bacteria may be of minimal significance under natural conditions, but can cause substantial problems when animals are crowded and stressed under culture conditions. The adverse effects associated with the use of antibiotics

include drug residue, bioaccumulation and resistance of pathogens, which threaten human consumers (Lee and Gao, 2012).

Justification for the Study

Catfish culture has contributed significantly to the growth of aquaculture in Nigeria on the potential effectiveness of some plant derived phyto constituents such as garlic extract that can serve as organic substance in treatment of fish disease. Its phytochemical constituents which are known to have antibacterial, anti-fungal, measure in treatment of *Clarias gariepinus* fingerlings. *Allium sativum* is a pungent herb and has been reported to inhibit bacterial growth, promote fish growth and enhancement of blood parameters (Metwally, 2009; Alam et al., 2016).

The use of organic product in recent years has been frequently researched. These products are often considered as easy to obtain and safe to use in terms consumption, such as garlic (Bilen et al., 2019; Elkordy et al., 2021).

Objectives of the Study

The objective of the study is to determine the phytochemical constituent of garlic extract (*Allium sativum*) and its influence on the treatment of bacterial and fungal infection in *Clarias gariepinus*.

Null Hypothesis

Null hypothesis: Garlic do not have significant content of photochemical for treating bacterial or fungal diseases ($p > 0.05$)

MATERIALS AND METHODS

Study Location

The experiment was conducted at the Microbiology Laboratory Department of Microbiology, Federal University of Technology Minna, Niger State, Nigeria and the Centre for Genetic Engineering and Biotechnology (CGEB), Federal University of Technology in Bosso Campus,

Processing of Plants Materials

Garlic bulbs were purchased from Kure market in Minna Niger State. One (1) kilogram of garlic bulb was used. The bulbs were peeled, washed under running tap water and pulverized. Using a pestle and mortar so as activate the various sulphur compound in garlic which was indicated by the pungent smell then it was further soak in a beakers containing 800ml of sterile water in a complete submersion.

Aqueous Extraction of Garlic

Garlic was extracted using cold maceration technique (Ahmad *et al.*, 2020). About 400g quantity of the pulverized *Allium sativum* bulbs were soaked with clean distilled water in a clean sterilized airtight container for three days (72 hours) at room temperature, while undergoing vigorous shaking at regular interval. The mixture was then filtered through muslin cloth and the concentrated filtrate was received in a sterile beaker. Using a water bath, the concentrated extract was subsequently transferred into clean sterile airtight glass container and stored in the refrigerator at 4 °C until required. The weight of the garlic extract was recorded and the percentage yield was estimated using El-Rokiek *et al.*, (2019) formula:

$$\%Yield = \frac{\text{Weight of dried extract}}{\text{Weight of plant sample}} \times 100 \dots\dots\dots(3.1)$$

Quantitative Determination of Phytochemicals

The quantitative phytochemical analysis of the obtained extract was carried out at the centre for genetic engineering and biotechnology (CGEB), Federal University of Technology Minna, Bosso campus.

Flavonoids determination

Ibrahim *et al.*, (2020) method was used to determine total flavonoid content of the extracts. of each extract was measured and added to a test tube containing 1.5 mL of absolute methanol, 0.1 mL of 10 % aluminum chloride, 0.1 mL of 1 % sodium acetate and 2.8 mL of distilled water. The tubes were incubated at ambient temperature for 30 minutes. The absorbance was read at 415 nm with double beam shimadzu UV spectrophotometer, UV-1800. Standard quercetin was used to prepare the calibration curve.

Phenol determination

Total phenol content of the extracts was determined using the method of Chaudhry *et al.*, (2022), 0.01 g of each extract was dissolved in 10 mL of distilled water, and 0.5 mL was oxidized by 2.5 mL of 10 % Folin-Ciocalteu's reagent which was then neutralized by 2 mL of 7.5 % sodium carbonate. Followed by vigorous shaking, the mixture was allowed to stand for 2 h. Finally, the absorbance was read at 765 nm using double beam Shimadzu UV spectrophotometer, UV-1800. Standard garlic acid was used to prepare the calibration curve.

Saponin determination

Saponin content of the extract was determined using the method of Adam *et al.*, (2023), 0.5 g of each extract was weighed and dissolved in 20 mL of 1N HCl and boiled in water bath at 80 °C for 4 h. The reaction mixture was cooled and filtered, 50 mL of petroleum ether was added and the ether layer was collected and evaporated to dryness. Thereafter, 5 mL of acetone-ethanol (1:1), 6 mL of ferrous sulphate and 2 mL of concentrated sulphuric acid were added and allowed to stand for 10minutes. The absorbance was taken at 490 nm. Standard saponins was used to prepare the calibration curve.

Alkaloid determination

Total alkaloid of the extracts was determined using method of Chaudhry *et al.* (2022), about, 0.5 g of each extract was weighed and dissolved in 5 mL of mixture of 96 % ethanol:20 % H₂SO₄ (1:1) and then filtered.1 mL of the filtrate was then added to a test tube containing 5 mL of 60 % H₂SO₄ and allowed to stand for 5 minutes. Thereafter, 5 mL of 0.5 % formaldehyde was added and allowed to stand at room temperature for 3 h. The absorbance was read at wavelength of 565 nm. Vincristine extinction coefficient (E₂₉₆, ethanol {ETOH} = 15136M⁻¹cm⁻¹) was used as reference alkaloid.

Tannin determination

Tannin content of the extracts was determined using the method of Palacios *et al.*, (2021), 0.2g of each extract was weighed into a 50 mL beaker and 20 mL of 50 % methanol was added to it and covered with para film and heated in water bath at 80°C for 1 hour. The reaction mixture was shaken thoroughly to ensure uniformity. The extract was then filtered into a 100 mL volumetric flask, and 20 mL of distilled water, 2.5 mL of Folin-Denis' reagent, and 10 mL of sodium carbonate were added and mixed properly. The reaction mixture was then allowed to stand for 20 minutes at room temperature for the development of bluish-green coloration. The absorbance was taken at 760 nm using double beam shimadzu UV-spectrophotometer, UV-1800. Standard tannic acid was used to prepare the calibration curve.

Data Analysis

Results were analysed statistically by using analysis of variance and Least Significant Difference test (LSD) according to the statistical system (SPSS-23). One-

Way ANOVA was used to compare anti-bacterial activities of Garlic extracts with Chlorophenicol. Excel was used to determine the yield of garlic extract

RESULTS AND DISCUSSION

RESULTS

Table 1: Yield of *Allium sativum* bulb extract

Extract	Weight(g)	Extract weight (g)	Yield (%)
<i>Allium sativum</i> bulb	400	37.69	9.422

Table 2: Phytochemicals Composition of *Allium sativum* bulb extract

Phytochemicals	Composition/concentration (mg/100 g)
Phenols	205.96
Flavonoids	62.85
Alkaloids	7.72
Tannins	9.39
Saponins	55.22

Table 3: Total bacterial and coliform count of infected *Clarias gariepinus* fingerlings

Infected <i>Clarias gariepinus</i>	Colony count (CFU/g)	
	TBC($\times 10^4$)	TCC($\times 10^3$)
1	10	23
2	6	9
3	26	1
4	16	8
5	23	6
6	27	1
7	11	12
8	20	4
9	12	7
10	9	1
11	30	16
12	20	1
13	14	6
14	37	12
15	22	9
16	17	2
17	16	7
18	7	7
19	24	18
20	13	5
21	8	11
22	25	9
23	37	14
24	16	7
25	21	9
26	9	19
27	26	7
28	35	3
29	14	20
30	11	16

Keys: TBC: Total bacterial count; TCC: Total coliform count; CFU: Colony forming unit;

DISCUSSION

Yield of crude extract of *Allium sativum* bulb

The yield of *Allium sativum* bulb extract is shown in table 4.1. The aqueous extract obtained weighed 37.69g and the estimated yield from 400g of *Allium sativum* bulb used for extraction was 9.4%. The *Allium sativum* bulb that yielded 37.69g (9.4%) is similar to the yield observed by Kallel et al. (2014) and Abd Elwahed et al. (2019). Kallel et al., (2014) reported a percentage yield of 26.5% using methanol and 50/50 methanol–water extracts while Abd Elwahed et al. (2019) reported 20% yield using bioregulators of two bio-regulators, indole acetic acid (IAA) and indole butyric acid (IBA). The observed discrepancy in result could be attributed to various factors majorly, the extraction method and quantity of extraction material used in each study. According to Alara et al. (2021), there are various extraction techniques including maceration, decoction, percolation, infusion, digestion, serial exhaustive extraction, and soxhlet extraction with each having its advantage and demerit on extraction outcome.

Quantitative phytochemicals screening of *Allium sativum* bulb extract

The phenol, tannin, flavonoid and alkaloid observed in garlic in this study were also observed by Gulfranz et al. (2014), Divya et al. (2017), Nazir and Chauhan (2019), Azizah et al., (2020) and Deepa and Sivakumar, (2023). Quantitatively, alkaloid (7.2%) was found to be the most abundant constituent, followed by tannin (4.80%), saponin (4.3%), flavonoids (2.18%) and phenols (0.80%). Their study also reported other phytochemicals including Glycosides, steroids, terpenoid and anthraquinones. However, Bar et al., (2022) did not observe alkaloids and flavonoid in garlic extract. The disparity observed in the phytochemical constituent of *Allium sativum* extract could be attributed to difference in solvents used in each study. According to Dzah et al., (2020) and Gil-Martín et al. (2022) the recovery of polyphenols from plant materials is influenced by the solubility of the phytochemical compounds in the solvent used for extraction.

Total bacteria and coliform count

The total bacteria and coliform count of infected *Clarias gariepinus* fingerlings is shown in Table 4.3. The total bacterial count of infected *Clarias gariepinus* ranged from 6×10^4 – 37×10^4 CFU/g with average count of 18.73×10^4 CFU/g the total coliform count ranged from 1×10^3 – 23×10^3 CFU/g with an average count of 9.00×10^3 CFU/g. In the current study the total bacterial count of infected *Clarias gariepinus* range from 6×10^4 – 37×10^4 CFU/g

with average count of 18.73×10^4 CFU/g while the coliform count ranged from 1×10^3 – 23×10^3 CFU/g with an average count of 9.00×10^3 CFU/g. Similar finding was reported by Afolabi et al. (2020) who assessed the bacterial loads of *Clarias gariepinus* obtained from cultured and natural habitats. They recorded a total bacterial count of 25.77×10^4 CFU/g. Higher Bacterial and coliform count was observed in the study of Ezeama et al. (2023), who reported a total bacterial count ($\times 10^{12}$ CFU/g) of 117.33 ± 11.554 while coliform count ($\times 10^{12}$ CFU/g) observed was 66.08 ± 5.51 . The observed microbial load in these studies could be as a result of exposure and consumption of bacteria for long time through food and water (Kebbi et al., 2020). The survival of these bacteria is dependent on the conditions prevailing in the aquatic environment and fish are often simply their hosts (Sehna et al., 2021). High bacterial abundance is not necessarily a disadvantage. If the bacteria are not pathogenic, as high bacterial abundance may indicate healthy organic matter recycling and re-mineralization (Kasozi et al., 2023).

Identification of bacterial isolates

Bacteria isolated from infected *Clarias gariepinus* include members of the genus *Lactobacillus*, *Staphylococcus*, *Bacillus*, *Micrococcus*, *Streptococcus*, *Enterobacter* and *Escherichia*. Represented by the species *Bacillus subtilis*, *Escherichia coli*, *Staphylococcus epidermidis*, *Micrococcus luteus*, *Enterobacter aerogenes*, *Staphylococcus aureus*, *Streptococcus* sp. and *Lactobacillus bulgaricus*. Similar findings were reported by Deborah et al., (2022) in their study on the occurrence of pathogenic bacteria associated with *Clarias gariepinus* in selected fish farms of Kumbotso Local Government Area of Kano State, Nigeria. From their study they reported the presence of *Escherichia coli*, *Micrococcus luteus*, *Enterobacter aerogenes*, *Pseudomonas aeruginosa*, *Staphylococcus epidermidis* and *Staphylococcus aureus*. Similarly, Ogbukagu et al., (2021) reported *Vibrio*, *Aeromonas*, *Pseudomonas*, *Lactobacillus*, *Staphylococcus*, *Microbacterium*, *Serratia*, *Proteus*, *Bacillus*, *Streptococcus*, *Citrobacter* and *Micrococcus*.

CONCLUSION

Eight bacteria which include; *Staphylococcus epidermidis*, *Bacillus subtilis*, *Streptococcus* sp., *Enterobacter aerogenes*, *Staphylococcus aureus*, *Escherichia coli*, *Micrococcus luteus*, *Lactobacillus bulgaricus* were isolated from infected *Clarias gariepinus* during the practical Phytochemical analysis

of the garlic extract reveal the presence of five secondary metabolite which includes phenols 205.96 (mg/100g), flavonoids 62.85 (mg/100g), alkaloids 7.72 (mg g/100g), Tannins 9.39 (mg/100g), saponins 55.22 (mg/100g).

Antibacterial activity of *Allium sativum* extract against suspected bacteria isolate tested show antibacterial activity. With highest activity observed at 1000mg/ml compared lower concentration of 800mg/ml, 600mg/ml, 400mg/ml respectively.

Garlic extract prove to be highly effective in treatment and control of bacteria disease infection in *Clarias gariepinus* at 1000mg/ml

RECOMMENDATIONS

- i. Further studies are needed to elucidate the Pathophysiological mechanism of action of garlic as well as its efficacy and safety in treatment of various diseases.
- ii. The extract mechanism of all ingredient and their long terms effect are not fully understood. Therefore studies should be carried out this area.
- iii. Further studies on the isolation and purification of active compounds present in *Allium sativum* bulbs should be carried out

REFERENCES

- Abd El-Hack, M. E., Samak, D. H., Noreldin, A. E., El-Naggar, K., & Abdo, M. (2018). Probiotics and plant-derived compounds as eco-friendly agents to inhibit microbial toxins in poultry feed: a comprehensive review. *Environmental Science and Pollution Research*, 25, 31971-31986.
- Abd Elwahed, M. S., Mahdy, H. A., El-Saeid, H. M., & Abouzienna, H. F. (2019). Effect of some bioregulators on yield quantity and quality of garlic plants (*Allium Sativum L.*). *Middle East Journal Applied Science*, 9(1), 17-24.
- Afolabi, O. J., Oladele, O. O., & Olususi, F. C. (2020). Assessment of bacterial loads of *Clarias gariepinus* (Burchell, 1822) obtained from cultured and natural habitats. *The Journal of Basic and Applied Zoology*, 81, 1-7.
- Ahmad, R., Ahmad, N., Riaz, M., Altarouti, M., Aloufi, F., AlDarwish, A., ...& Khan, Z. (2020). Optimization of extraction and quantification technique for phenolics content of garlic (*Allium sativum*): An application for comparative phytochemical evaluation based on cultivar origin. *Biomedical Chromatography*, 34(11), e4942.
- Ayele, T. A. (2015). Growth performance and survival rate of African catfish larvae *Clarias gariepinus* (Burchell 1822) fed on different types of live and formulated feeds. *MSc, University of Natural Resources and Life Science, Vienna, Austria.*
- Azizah, Z., Yani, P., & Yetti, R. D. (2020). Antioxidant activity ethanol extract of garlic (*Allium sativum L.*) and black garlic. *Int. J. Res. Rev*, 7(9), 94-103.
- Ben, Y., Fu, C., Hu, M., Liu, L., Wong, M. H., & Zheng, C. (2019). Human health risk assessment of antibiotic resistance associated with antibiotic residues in the environment: A review. *Environmental research*, 169, 483-493.
- Bravo, S. (2012). Environmental impacts and management of veterinary medicines in aquaculture: the case of salmon aquaculture in Chile. *Improving biosecurity through prudent and responsible use of veterinary medicines in aquatic food production*, 547, 11-24.
- Choudhary, S., Noor, M. U., Hussain, M. S., Mishra, M., & Tyagi, S. (2022). Pharmacological properties and phytoconstituents of garlic (*Allium sativum L.*): A review. *Biological Sciences*, 2(4), 338-346.
- Chaudhry, F., Ahmad, M. L., Hayat, Z., Ranjha, M. M. A. N., Chaudhry, K., Elboughdiri, N., ...& Uddin, J. (2022). Extraction and evaluation of the antimicrobial activity of polyphenols from banana peels employing different extraction techniques. *Separations*, 9(7), 165.
- Deborah, A. A., Saidu, L., Sonnie, O., Adakole, A. S., Susan, D. M., & Omodolapo, O. M. (2022). Antibiotic resistance patterns of bacteria isolated from *Clarias gariepinus* farms in Kaduna state, Nigeria. *Iranian Journal of Veterinary Science and Technology*, 14(1), 29-38.
- Deepa, B., & Sivakumar, T. (2023). Screening of Phytochemicals and in vitro studies of Garlic: An Updated review. *International Journal of Engineering Technology and Management Sciences*, 7, 1-2.
- Elkordy, A. A., Haj-Ahmad, R. R., Awaad, A. S., & Zaki, R. M. (2021). An overview on natural product drug formulations from conventional medicines to nanomedicines: Past, present and future. *Journal of Drug Delivery Science and Technology*, 63, 102459.
- Fetsch, A., & Jöhler, S. (2018). Staphylococcus aureus as a foodborne pathogen. *Current Clinical Microbiology Reports*, 5, 88-96.
- Gajewska, J., & Chajęcka-Wierzchowska, W. (2020). Biofilm formation ability and presence of adhesion genes among coagulase-negative and coagulase-positive staphylococci isolates from raw cow's milk. *Pathogens*, 9(8), 654.

Gulfraz, M., Imran, M., Khadam, S., Ahmed, D., Asad, M. J., Abassi, K. S., ...& Mehmood, S. (2014). A comparative study of antimicrobial and antioxidant activities of garlic (*Allium sativum* L.) extracts in various localities of Pakistan. *African Journal Plant Sciences*, *8*, 298-306.

Hajar-Azhari, S., Abd Rahim, M. H., Sarbini, S. R., Muhiadin, B. J., Olusegun, L., & Saari, N. (2021). Enzymatically synthesised fructooligosaccharides from sugarcane syrup modulate the composition and short-chain fatty acid production of the human intestinal microbiota. *Food Research International*, *149*, 110677.

Hamed Aboelkhair, H. A. (2022). *Study of Microbial Enhanced Oil Recovery using Indigenous Bacteria in Egyptian Oil Reservoirs* (Doctoral dissertation, London South Bank University)

