

Full Length Research Paper

Spatial Distribution of Nitrate in the Drinking Water Sources Found in Ethiopia; *Retrospective Study*

^{*1}Andualem Mekonnen, ²Tsigereda Assefa, ²Kuribel Tesfaye, ²Derbu Getahun

¹Center for Environmental Science, College of Natural Science, Addis Ababa University, P.O.Box 33348, Addis Ababa, Ethiopia

²Ethiopian Health and Nutrition Research Institute, Non-Infectious Diseases Research Team

*E-mail: andumk21@gmail.com, andualem.mekonnen@aau.edu.et

Accepted 9th, December, 2014

This retrospective study aimed to investigate the spatial variation of nitrate level in the drinking water sources found in Ethiopia. In the study, the Ethiopian Health and Nutrition Research Institute water quality database from the year 1993 to 2007 was used. The results of the water sample indicated that nitrate concentration were varied from below 0.00mg/l to 1295.00mg/l. Nationally, 15.3 % (n=186) of the wells, 10 % (n = 33) of the springs and 12.4 % (n=21) of the taps had nitrate concentration exceeding the threshold value of 20mg/l and 5.7 % (n = 70) of well water and 2.7 % (n = 9) of spring water samples had nitrate concentration higher than 50mg/l. The average nitrate concentration in the well waters exceeding the WHO guideline was observed in Dire Dewa (104.8mg/l) and exceeding the threshold value were found in Somali (37.0 mg/l), Afar (34.9 mg/l), Harari (26.3 mg/l) and Addis Ababa Regions (20.5mg/l). Nitrate in all the spring water was below the threshold value except Dire Dewa and Harari regions. In those water sources with high nitrate concentration, management action should be taken to protect the water from further contamination.

Keywords: Nitrate, drinking water, well water, spring water, tap water

INTRODUCTION

Drinking water containing high level of nitrate is becoming public health concern due to endogenous reduction to highly toxic compounds such as; nitrites and nitrosamine (Ward et al., 1994; Chiu and Tsai, 2007). Methemoglobinemia is the primary adverse health effect associated with human exposure to nitrate or nitrite (Deana *et al*, 2010). Infants less than six months old are the primary vulnerable group to acquired methemoglobinemia from exposure to nitrates in drinking water (Deana *et al*, 2010; Ayebo et al., 1997). Pregnant women are the other vulnerable group affected by high level nitrate exposure in drinking water (Deana et al, 2010). The second adverse health effect of nitrates in drinking water is the possibility of increased cancer risk (via the bacterial production of N-nitroso compounds), hypertension, increased infant mortality, central nervous system birth defects, diabetes, spontaneous abortions respiratory tract infections and changes to the immune system (Gheisari et al., 2005; Babiker, 2004; Gupta et al., 2000; Ward et al., 1994; Chiu and Tsai, 2007).

Groundwater with nitrate concentration exceeding the threshold of 20 mg/L is considered contaminated as result of human activities (Spalding and Exner, 1993). Recent studies revealed that water contamination with nitrate is globally growing problem due to the high rate of population growth and increasing consumption (Jalali, 2005). In most European countries, nitrate levels in rivers and ground waters have increased gradually over the last decade mainly as a consequence of large-scale agricultural application of manure and fertilizers, thereby threatening drinking water quality (Stringer, 1988). World Health Organization (WHO) guidelines on the quality of drinking water indicate a maximally admissible nitrate concentration of 50.0 mg nitrate/L (Stringer, 1988; Duijvenbooden, 1989).

In Ethiopia, well water and spring water are the major water supply source used in both urban and rural areas (Gebrekidan and Samuel, 2011). The results of some fragment studies indicated that there is an increasing problem of water pollution. This could be due to



Figure 1: Map of Ethiopia and the nine regional states

contamination resulted from soil erosion, agricultural runoff, domestic and industrial wastes. However, there is no comprehensive information about the water quality status of the country. Therefore, this paper aims to examine the concentration of nitrate in the well water, spring water and tap water which were collected and analysed from different regions of Ethiopia.

MATERIAL AND METHODS

Country description

Ethiopia found in the Horn of Africa and located between 3°N and 15°N of the equator and 33°E and 48°E longitudes. The country is surrounded by Sudan in the west, Kenya in the south, Somalia in the southeast, Djibouti in the east and Eritrea in the north and northeast. The total area of the country is about 1,127,127 km². In the country, there is great geographical variation, with a topography ranging from 4550 metres above sea level to 110 metres below.

There are nine administrative nine regions as indicated in figure 1. These regions are Afar, Amhara, Benishangul-Gumuz, Gambella, Harari, Oromia, Somali, Tigray and the Southern Nations, Nationalities and Peoples Region (SNNPR) and two administrative

councils (Addis Ababa and Dire Dawa). Each regional state is subdivided into zones and *woredas*.

Sample collection and analysis:

This retrospective study of nitrate was conducted using the Ethiopian Health and Nutrition Research Institute water quality database. The water samples were collected by Environmental health professional using plastic container. The nitrate concentration in the water sample was measured using the phenoldisulfonic acid method according to Standard Method of Water and Wastewater analysis (APHA, 1965). Nitrate test results data between 1993 to 2007 organized and analyzed using excel. The total numbers of the water samples assessed were 1719 and from this 1220 were from well water, 330 from spring and the remaining 169 were from tap water. The concentration of nitrate in the water samples were compared with WHO guideline.

RESULTS AND DISCUSSION

Nitrate concentration in the water samples varied from less than 0.00mg/l to the maximum of 1295.00mg/l throughout the country. The samples were classified into three groups based on their nitrate concentrations.




Figure 2a: Number of samples at different levels of nitrate b) Percentage of samples at different levels of nitrate

These included the following; low (<20 mg/L), medium (≥20 to <50 mg/L) and high (>50 mg/L) (Figure 2a and b).

Nitrate concentration in the lower class indicates samples with a low risk for human or environment. The medium class involves samples with high nitrate concentration that indicate the impact of human activities in those water sources (Spalding and Exner, 1993; Mutewekil et al., 2008). About 15.3 % of the wells water samples (n=186), 10 % of the springs water samples (n = 33) and 12.4 % of the tap water samples (n=21) had nitrate concentration higher than the threshold value (20mg/L). The water samples in the higher class contain nitrate concentration that exceeded the WHO drinking water recommendation guideline (WHO, 1993). 5.7 % of well water samples (n = 70), 2.7 % of spring water samples (n = 9) and 2.4 % of the tap water samples (n = 4) had nitrate concentrations higher than the maximum allowable nitrate concentration (50mg/l). This information is further broken down by region (Table 1 and 2). Moreover, the areas which have high nitrate concentration above the national and WHO standard are summarized in Table 3.

In Afar region, 106 well water, 8 spring and 7 tap water samples were collected and analyzed. The maximum and average nitrate concentrations detected were 294.5 and 34.9, 2.3 and 2.66, and 6.7 and 17.7 mg/L as NO_3^- , for well, spring and tape water, respectively (Table 1 and 2). Nitrate concentration in the spring and tape water samples was very low. Nitrate concentration in all the spring and tape water samples were complying with the national and WHO standard of nitrate. However, the average concentration of nitrate in all the well water samples was above the threshold value of nitrate. Moreover, 19.8% of the total samples (n=21) had nitrate

concentration above the threshold value (20mg/L) and 10.4 % of samples (n=11) were above the national standard. The nitrite concentration of 6.6% of the sample (n=7) was also above 3mg/l which is the national and WHO standard of nitrite (3mg/l). Water samples, which had high concentration of nitrate, were collected from Dubty, Assaita, Middle Awash, Elidar, Afambo, Hamiltola, Handeg, Geyreni and Gilifega village (Table 3). All these villages are located in the lower valley of the Awash River. In the Lower Awash Valley, salinity problems and evapotranspiration are very high throughout the area (Tadesse et al., *undated*). In Dupty, the potential evapotranspiration (PET) is estimated to be ten times the average annual rainfall (Tadesse et al., *undated*). Study indicated that in the saline groundwater, evaporation may increase nitrate concentration in the ground water (Piomleart, 1995). This could be one of possible factors for high concentration of nitrate. The other possible factors could be the intensive agricultural activities in the region. In the lower valley, large scale irrigated farming is common used. More than 62500 hectare of land is cultivated using irrigation (Tadesse et al., *undated*).

In Addis Ababa, 189 well water, 11 spring water and 65 tape water samples were collected from all part of the city. The concentration of nitrate in the tape water were very low in all the samples with an average value of 1.6mg/L and nitrate in all the sample were with in the national standard. The mean value and maximum value of spring water samples were 12.3mg/l and 69.6 mg/l, respectively. One spring water sample only did not comply with the standard. However, nitrate concentrations were high in the well water samples. The average value of nitrate concentration was 20.5mg/l, which is above the threshold value, and maximum value

Table 1: Nitrate and nitrite concentration in well waters collected from different regions

Region	Total *No.	Nitrate concentration (mg/L)								Nitrite	
		≥20 - <50		>50mg/L		≤0.00mg/L		Mean value	Max. value	>3mg/L	
		*No.	(%)	*No.	(%)	*No.				*No.	(%)
Afar	106	10	9.4	11	10.4	12	11.3	34.9	294.5	7	6.6
Addis Ababa	189	33	17.4	20	10.6	20	10.6	20.5	327.3	3	1.5
Amhara	121	9	7.4	8	6.6	12	9.9	11.3	166	3	2.4
Benishangul	46	0	0	3	6.5	21	45.7	14.2	409	0	0
Dire Dawa	19	4	21	12	70.6	-	-	104.8	271.6	0	0
Gambela	12	1	8.3	0	0	6	50	3.4	40	0	0
Oromia	568	36	6.3	6	1.1	119	20.9	6.09	114.3	8	1.4
SNNPR	54	0	0	0	0	17	31.5	2.72	1.41	0	0
Somali	22	2	9.1	4	18.2	6	27.3	37	376.6	2	9
Tigray	72	17	23.6	4	5.6	9	12.5	19	235	7	9.7
Harari	12	4	33.3	2	16.7	1	8.3	26.3	79.14	0	0

*No. = Number of samples

Table 2: Nitrate concentration in spring and tape water collected from different regions

Region	Nitrate concentration in spring water						Nitrate conc. in tape water			
	Total *No.	No. sample			Mean value (mg/l)	Max. value (mg/l)	Total *No.	*No. > 50 (mg/l)	Mean value (mg/l)	Max. value (mg/l)
		≥20 - <50 (mg/l)	>50 (mg/l)	≤0.00 (mg/L)						
Afar	8	0	0	1	2.3	2.66	7	0	6.7	17.7
Addis Ababa	11	1	0	2	12.3	69.6	65	0	1.6	4.9
Amhara	57	7	3	11	12.6	98.4	3	0	0.3	0.9
Benishangul	3	0	0	1	0.24	0.36	0	-	-	-
Dire Dawa	4	2	1	1	49.4	126.8	0	-	-	-
Oromia	126	7	1	30	6.5	314.5	48	0	3.6	26.4
SNNPR	40	1	1	17	5.1	61	5	0	2.1	5.12
Tigray	19	5	0	2	14	38.3	4	0	0.7	2.2
Harari	4	0	1	-	58.4	260	5	0	21.3	36.7

*No. = Number of samples

Table 3: Woreda/village which have high concentration of Nitrate in their water sources

Region	Zone	Woreda/ village	Sample source	NO ₃ ⁻ (mg/l)
Afar	Zone-1:	Dubty, Asayita, Middle Awash, Elidar, Afambo, Hamiltole, Handeg, Geyreni and Gilifega	well	256.4
Amhara	N/Gonder	Debremariam spring/Azezo	spring	51.6
	N/ Shewa	Eyerusalem, Gozie, Kecheni	well	83.3
	Wagimra (Sekota)	Kative, Kitibe Beleka, Ekmetseskue, Selessia, Tsebiyo cherkose	Spring well	77 92.4
Benishanguel	Dabatie (pawie)		well	193.6
Oromia	E/ Shewa:	Metahara, Debrezeit, and Akaki	well	72.9
Somali	Jijiga:	Jijiga, Jerere valley (KebriBeyah)	well	97.4
Tigraye	Kilte-Awlaelo (E/zone),	Sheraso town, Chekentemy, Mai Daga / Axum, Mekele	well	112.8

detected was 327.3 mg/L. Out of 189 well samples, 28% of the samples (n=53) were above the threshold and nitrate concentration in 10.6 % of the samples (n=20) exceed the national standard. The high nitrate levels are likely caused by a lack of proper sewers and other waste disposal facility. The majority of the households (75%) in Addis make use of a pit latrines discharging to open drains and about 15% have flush toilets and septic tanks, these likewise often discharging to open drains (Alemayehu et al., 2009; AAWSSA, 2008). According to Yates (1985) improperly used septic tanks could be the major contributors of waste to the subsurface water. On the other hand, twenty samples (10.6%) had nitrate concentration below 0.00mg/l and all these samples were collected from the foot of Entoto Mountains. Nitrate contamination Addis Ababa aquifer is well documented in different studies (Taye, 1998; UN, 1989).

One hundred twenty one well water and fifty seven spring water samples were collected and analyzed from different parts of Amhara region. The average and maximum value of nitrate concentration were 11.3 and 166mg/L for well water, and 12.6 and 98.4 mg/l for spring water (Table 1 and 2). Nitrate concentrations above the national standard were detected in 6.6 % of the well water samples (n=8) and 5.2 % of the spring water samples (n=3). Nitrite concentration was also high in 2.4 % of well samples (n = 3). Nitrate concentration above the national standards was detected in some areas of three zone in Amhara region (Table 3). In Wagimra zone, spring water from Kative and Kitibe, and well water from Beleka, Ekmetsaku Selessia and Tsebiyo cherkose had nitrate concentrations which exceed 50mg/L. One spring water from North Gonder (Azezo) had also concentration of nitrate above the standard. On the other hand, in North Shewa zone only well water from Eyerusalem, Gozie and Kecheni had nitrate concentration above the standard (Table 3).

In Benishangul region, 46 well water and 54-spring water samples were analyzed. The average and maximum value of nitrate detected were 14.2 and 409, and 0.24 and 0.36mg/l for the well and spring, respectively (Table 1 and 2). Twenty one samples (45.7 %) had nitrate concentration below 0.00mg/L. On the other hand, water samples from three wells (6.5%) exceeded the maximum nitrate contamination levels allowed in drinking water. These water samples were collected from Metekel zone at Pawie (Table 3).

The numbers of well, spring and tape water samples collected from different parts of the Oromia region were 568, 126 and 48, respectively (Table 1 and 2). The average and maximum value of nitrate concentration in the samples were 6.09 and 114.3 mg/l for well water, 6.5 and 314 mg/l for spring water, and 3.6 and 26.4 mg/L for tape water samples, respectively (Table 1 and 2). Out of 568 well samples, 20.9% of samples (n=119) and 23.8% of spring samples (n=30) had nitrate concentration below 0.00mg/L. On the other hand, nitrate

concentration in 0.8% of the spring and 1.1% of the well samples (n = 6) were above the national standard. From the well water samples, 1.4 % of the samples (n=8) had also nitrite concentration above 3mg/L (above the standard). Well waters from Metehara, Akaki and Debreziet had average nitrate concentration of 72.9 mg/L which is above the acceptable level (Table 3). The high concentration of nitrate in East Shewa zone could be attributed to both agricultural practice and sewage discharge.

In Dire Dewa region, 19 well water and 4 spring water samples were analyzed. The average and maximum value of nitrate concentrations detected were 104.8 and 271.6mg/l, and 49.4 and 126.8 mg/l, for well and spring water, respectively (Table 1 and 2). Water samples from 18 of these wells (94.7%) had nitrate concentration above the threshold value of nitrate. Nitrate concentration in 12 well samples (70.6%) and 1 spring sample exceed the WHO and national standard. Nitrate concentration in the well water samples collected from Harari region were in the range of 0.00 to 79.14 mg/L. The average nitrate concentration was 26.3mg/l which is above the threshold value for nitrate. Out of the total samples, two samples (16.7%) had nitrate concentration that exceeds 50mg/L, the maximum acceptable level.

Twenty two and twelve well water samples were analyzed from Somali and Gambela region, respectively. In Somali region, the average nitrate concentration was 37mg/L which is above the maximum threshold value and the maximum value detected was 376.6mg/L. Nitrate concentration in 18.2% of the sample (n=4) exceed the national standard. The high value of nitrate was associated with human actives. However, in Gambela region, both the average value (3.46mg/L) and the maximum value (40mg/L) were within the national standard. The high concentration of nitrate in the groundwater resource of Dire Dewa, Harari and Somali region could be caused by factors similar to those causing nitrate problems in Addis Ababa.

In SNNRP region, 54-well water, 40-spring water and 5-tape water samples were analyzed. The concentration of nitrate in water samples from all wells were detected very low which had an average value of 1.41mg/L and a maximum value of 2.72 mg/L. One sample from the spring water had nitrate concentrations which exceed the maximum allowed levels for drinking water.

In Tigray region, 72 well water, 19 spring and 4 tape water samples were analyzed. The average and maximum nitrate concentrations detected were 19 and 235, 14 and 38.3, and 0.7 and 2.2 mg/L as NO_3^- , for well, spring and tape water, respectively (Table 1 and 2). All spring and tape water samples were comply with national standard of nitrate. However, nitrate concentrations in 21 well water samples exceed the threshold value of nitrate and 4 samples were above the national standard. The high levels nitrate were recorded in the water sample collected from Kilt-Awlaelo(E/zone), Sheraso town, Chekentemy, Mai

Daga / Axum and Mekele (Table 3). Tekle *et al* (2004) assessed nitrate pollution in Aynalem and Mekele groundwater sources of drinking water for Mekelle city. It was reported that nitrate levels in the downstream wells of Mekele had above 50mg/l. The reported possible sources of nitrate are animal feedlots and municipal wastes. NEDECO (1998) had also reported high nitrate concentrations in polluted wells around large towns such as Mekele, Indasilase and Shiraro (NEDCO, 1998).

CONCLUSION

The result of this study indicated that nitrate concentration in the water samples varied from below 0.00mg/l to 1295.00mg/l. Nationally, 15.3 % (n=186) of the wells, 10 % (n = 33) of the springs and 12.4 % (n=21) of the taps water samples had nitrate concentration exceeding the threshold value (20mg/l). In addition, 2.7 % (n = 9) of spring water samples and 5.7 % (n = 70) of well water samples had nitrate concentrations higher than 50mg/l. The highest concentration of nitrate were found in the water samples collected from Dire Dewa, Harari, Afar and some water samples collected from Addis Ababa, Amhara, Oromia, Tigary and Somali regions. On the other hand, the highest compliance was recorded in SNNRP, Gambela and Benishangul region. The highest concentration of nitrate in the water samples could be as a result of contamination from agricultural runoff, refuse dump, human and animal waste. In those water sources with high nitrate concentration, the level of nitrate should be reduced to the acceptable level before the water being used for domestic purpose. Moreover, water resource management action should also be implemented to protect the water resources from getting contamination.

REFERENCE

- AAWSSA (2008). Water production and distribution in Addis Ababa. Addis Ababa Water Supply and Sewerage Authority, Addis Ababa.
- Abdulrahman I. A., Abdullah M. A., Abdullah I. A. and Khan A. (2010). Assessment of nitrate concentration in groundwater in Saudi Arabia. *Environ Monit Assess* 161:1–9
- APHA (1965). Standard methods for the examination of water and wastewater, 12th Ed. Am. Pub. Health Assoc., New York, p. 195-98
- Ayebo A, Kross B, Vlad M, Sinca A (1997). Infant methemoglobinemia in the Transylvania region of Romania. *Int J Occup Environ Health* 3(1):20-29
- Babiker IS, Mohamed MAA, Terao H, Kato K, Ohta K (2004). Assessment of groundwater contamination by nitrate leaching from intensive vegetable cultivation using geographical information system. *Environment International* 29(8): 1009-1017
- Chiu HF, Tsai SS (2007). Nitrate in Drinking Water and Risk of Death from Bladder Cancer: An Ecological Case-Control Study in Taiwan. *J. Toxicol. Environ. Health, Part A* 70:1000-1004
- Criss RE, Davisson ML (2004). Fertilizers, water quality, and human health. *Environmental Health Perspectives* 112(10): A536-A536
- Deana MM, Lorraine CB, Rita M, Lora EF, Barbara L, Carolyn PM (2010). Nitrates in drinking water and methemoglobin levels in pregnancy. *Environmental Health* 9:60
- Duijvenbooden W, Matthijsen JM (1989). Integrated Criteria Document Nitrate. National Institute of Public Health and Environmental Protection Report no. 758473012. Bilthoven, The Netherlands,
- ECETOC (1988). Nitrate and drinking water. Brussels, European Chemical Industry Ecology and Toxicology Centre (Technical Report No. 27).
- Fields S (2004). Global nitrogen: Cycling out of control. *Environ. Health Perspect* 112:A557 A563.
- Gebrekidan Mebrahtu, Samuel Zerabruk (2011). Concentration of Heavy Metals in Drinking Water from Urban Areas of the Tigray Region, Northern Ethiopia. (*MEJS*) 3 (1):105-121
- Gheisari MM., Messripour M, Hoodaji M, Noroozi M, Abdollahi A (2005). Nitrate Intake from Drinking Water in Isfahan in 2004. *J. Sci., Islamic Republic of Iran* 16(2): 113-116
- Gupta SK, Gupta RC, Gupta AB, Seth AK, Bassin JK, Gupta A (2000). Recurrent acute respiratory infections in areas with high nitrate concentrations in drinking water. *Environ Health Perspect* 108(4):363-366.
- Jacks G, Sharma VP (1983). Nitrogen circulation and nitrate in ground water in an agricultural catchment in southern India. *Environmental geology* 5(2):61-64.
- Jalali M (2005). Nitrates leaching from agricultural land in Hamadan, Western Iran: Agriculture, Ecosystems and Environment 110
- Mutewekil MO, Fayez YA, Nezar AH, Adnan MM, Faisal SA (2008). Assessment of nitrate contamination of karst springs, Bani Kanana, northern Jordan *Revista Mexicana de Ciencias Geológicas*, v. 25, núm. 3, p. 426-437
- NEDECO (1998). Tekeze River Basin Integrated Development Master Plan Project, Executive Summary, Netherlands Engineering Consultants. Addis Ababa.
- Piromleart S (1995). Nitrate Affected Ground water in Northeast Thailand. International conference on Geology, Geotechnology and Mineral Resources of Indochina. Khan Kaen, Thailand.
- Spalding RF, Exner ME (1993). Occurrence of nitrate in groundwater –a review: *J. Environ. Qual.* 22: 392-402.
- Stringer DA (1988). Nitrate and Drinking Water. European Chemical Industry Ecology and Toxicology Centre Technical Report no. 27, Brussels, Belgium
- Syed SK, Saadat AK (2005) Level of Nitrate and Nitrite Contents in Drinking Water of Selected Samples Received at Afpngmi Rawalpindi. *Pak J Physiol* 1(1-2)
- Tadesse G, Sonder K, Peden D (2009). *undated*. The Water of the Awash Basin. A future challenge to Ethiopia. Last accessed on 17 th of Feb. 2009 at:
- Tamiru Alemayehu (2001). The causes for the surface and groundwater quality deterioration in Addis Ababa. 4th Theoretical Chemistry workshop in Africa (TCWA). Addis Ababa, Ethiopia.
- Taye A (1999). Pollution of the hydrogeologic system of Dire Dawa. Preprint for the 25th WEDC Conference on Integrated Development for Water Supply and Sanitation. Addis Ababa, Berhanena Selam Printing Enterprise.

- Tekele Kelali A, Yoshida I, Harada M. (2004). Nitrate Concentration in Drinking Groundwater Wells of Mekelle, Ethiopia (I) *J Rainwater Catchment Syst* 10 (1):1-5
- UN (1989). Ethiopia In: Ground water in Eastern Central and Southern Africa. Natural Resource/water series No.19, United Nations, New York, PP 84-95
- Wakida TF, Lerner DN (2005). Non-agricultural sources of groundwater nitrate: a review and case study: *Water Res.* 39: 3-16.
- Ward MH, DeKok TM, Levallois P (2005). Workgroup report: drinking-water nitrate and health-recent findings and research needs. *Environmental Health Perspectives* 113:1607-1614.
- WHO (1993). Guidelines for drinking water quality recommendations: Geneva, World Health Organization, 2nd edition.
- WHO. (1996). Guidelines for Drinking Water Quality (Vol. 2) (2nd edn.). World Health Organization, Geneva, Switzerland.
- Yates MV (1985) Septic-tanks density and groundwater contamination: *Groundwater*, 23, 586-591